Lake Wise Info Sheet



Shoreland Best Management Practices for Lake-friendly Living.

Benefits









Visual Appeal

Low Cost

Low Maintenance

Protection & Resiliency

Many BMPs for managing ticks are also VT DEC suggested BMPs for shorelands

Related Info Sheets:

Lakeshore Buffers
Planning Pathways

Restore Natural Plant Communities

TICKS & SHORELANDS

Minimize risk & protect water quality


Description.

Manage for native vegetation and pathways along shorelands to minimize risk of human contact with ticks and prevent tickborne diseases, including Lyme Disease and Anaplasmosis.



Native plant communities help to minimize tick populations.

It is important to understand the tick life cycle and conditions that cause tick populations to increase in order manage your property to reduce risk of contact.



How ticks spread.

Applicability.

Deer and mice are common hosts to ticks and help them to spread. Mice act as hosts for ticks in their larvae stage and deer are hosts to nymph and adult ticks.

^Tick life cycle.

Strategies to minimize contact with ticks on lakeshores.

1. Plant and promote native species.

Native plants diversify habitat for birds and other wildlife. Birds, such as owls, eat ticks in addition to other bugs and mice, lowering the tick population.

Native plant communities create an environment that is less protective of ticks and mice due to varied plant heights and growth forms. Woodland environments maintain a humidity level that does not necessarily promote tick growth.

Lyme disease.

Lyme disease is the most commonly reported tickborne disease in Vermont. It is caused by the bacteria *Borrelia burgdorferi* and is spread by the bite of a blacklegged tick. Ticks become "infected" during their larval stage when they first feed on blood of animals.

Lake Wise Info Sheet

TICKS & SHORELANDS



2. Plant deer resistant native species.

Planting native species that deer do not typically browse on helps to prevent the spread of ticks nearby. The following is a limited selection of native plants that are resistant to deer browsing:

Shrubs Native dogwoods Winterberry Holly Summersweet Fragrant Sumac **Viburnums**

Bayberry Shrubby Cinquefoil Witchhazel

X Trees

Birches Pines Junipers/ E. Red Cedar **Spruces** Maples Serviceberries Elderberries

Herbaceous Plants

Most ferns Columbine Most grasses Echinacea Anise Hyssop Lobelias **Asclepias** Liatris Asters Lupine Baptisia Rudbeckia Blue Flag Iris Wintergreen Bee Balm/Bergamot Yarrow

Not all ticks spread disease.

There are 15 different types of ticks in Vermont, but only six are known to bite humans and spread diseases. However, 99% of tickborne diseases reported to the Vermont Department of Health are caused by the blacklegged tick.



Lone Star Tick (Amblyomma americanum)







NOTE: Relative sizes of several ticks at different life stages

Engorged female Ixodes scapularis tick. Color may vary [^]Ticks that commonly bite humans.

Prevent tick bites with repellents registered by the EPA, like DEET (note: DEET should not be used on infants), Picaridin, and Oil of Lemon Eucalyptus. Permethrin can be used on clothing, but not skin. And always check yourself, children, and pets for ticks after spending time outdoors.

3. Maintain lake access pathways.

Defined mown or mulched pathways minimize the opportunities for brushing against ticks while still allowing natural vegetation to grow along the shore. Under the Shoreland Protection Act, one 6-foot-wide pathway is allowed without a permit.

4. Remove invasive plants.

Research studies have indicated that some invasive plant species create better conditions for ticks to thrive than healthy native forest ecosystems. University of Connecticut researchers found that invasive Japanese Barberry (Berberis thunbergia) increases the occurrence of Lyme disease-infected ticks fourfold because it creates the perfect, humid environment for ticks and protection for mice to hide in the dense branches.

For more information...

Healthvermont.gov/diseasecontrol/tickborne-diseases





